

Modulation of Innate and Adaptive Immune Responses by Arabinoxylans

Abdulmannan Fadel^{*1,7}, Andrew Plunkett¹, Weili Li², Vivian Elewosi Tessu Gyamfi¹,
Rosemarie Roma Nyaranga¹, Fatma Fadel³, Suaad Dakak⁴, Yazan Ranneh⁵, Yasser Salmon⁶,
and Jason J Ashworth⁷

^{*1} Department of Food and Nutrition, School of Health Psychology and Social Care,
Manchester Metropolitan University, Manchester M15 6BH, United Kingdom

² Institute of Food Science & Innovation, University of Chester, Chester CH1 4BJ, United
Kingdom

³ Independent Researcher, Al-Baha University, Saudi Arabia

⁴ Faculty of Pharmacy and Medical Sciences, Al-Ahliyya Amman University, Jordan

⁵ Department of Nutrition and Dietetics, Faculty of Medicine and Health Sciences, Universiti
Putra Malaysia, Serdang, Malaysia

⁶ Veteriner Fakultesi, Istanbul Universitesi, Turkey

⁷ School of Healthcare Science, Faculty of Science and Engineering, Manchester
Metropolitan Univeristy, Manchester M1 5GD, United Kingdom

Corresponding author: Dr. Abdulmannan Fadel
Email: man_fadel@hotmail.com

24

25

26

Modulation of Innate and Adaptive Immune Responses by Arabinoxylans

Abstract

Humans are exposed to harmful pathogens and a wide range of noxious substances every day. The immune system reacts to, and destroys, these pathogens and harmful substances. The immune system is composed of innate and adaptive immunity, which liaise to protect the host and maintain health. Foods, especially cereals, have been reported to modulate the immune response. Arabinoxylans are non-starch polysaccharides that have been shown to possess immune-modulatory activities. This review article discusses the fundamentals of the immune system and provides an overview of the immunomodulatory potential of arabinoxylans in conjunction with their structural characteristics and proposed similarities with lipopolysaccharides.

Practical applications

Understanding how the immune system works is of vital importance to prevent unnecessary or excessive inflammatory responses. Consumption of arabinoxylans has been shown to possess immunomodulatory potential. However, their mechanism of action has not been elucidated. Arabinoxylans share some similarities with lipopolysaccharides (LPS), a molecule that induces substantial and sometimes excessive immune responses such as fever following infection by pathogens. Thus, we propose that arabinoxylans might possibly act on the same receptor as LPS. Competition between dietary-derived arabinoxylans and LPS at a shared receptor would then have the potential to inhibit or attenuate excessive LPS-induced inflammatory responses that are typical of infection/fever. In the absence of infection and consequently no competition at the LPS receptor, consumption of dietary arabinoxylans may protect against the risk of

infection by moderately activating the receptor and heightening natural (background) levels of immunity.

Keywords: Arabinoxylans, lipopolysaccharides, innate immunity, adaptive immunity, nitric oxide, Toll-like receptor

1. Introduction

The human body is continually exposed to pathogens or harmful agents. The immune system possesses a set of defence mechanisms against these harmful pathogens and is composed of innate (non-specific) and adaptive (specific) immunity (Nicholson, 2016). The innate immunity encompasses chemical, microbiological and physical barriers, in addition to other elements of the immune system such as monocytes, macrophages, neutrophils, the complement system and cytokines (Sperandio et al., 2015). The innate response is immediate and non-specific, unlike the adaptive immunity, which is considered the hallmark of the immune system with its specific, yet slower response (Iwasaki & Pillai, 2014; Iwasaki & Medzhitov, 2015). Mounting an immune response against harmless foreign molecules is unnecessary and can lead to fatal outcomes such as anaphylactic shock (Patosuo, 2014). This deleterious response is typically avoided because the adaptive immune response is triggered by the innate immune system only when the latter recognises molecules of an attacking pathogen (Bonneaud et al., 2003; Lu et al., 2016).

Performance of the immune system is vital for defending the body from pathogens and it plays a crucial role in health homeostasis (Nairz et al., 2013). It has been suggested that ingestion of foods with immune-modulatory effects is able to prevent deterioration of immune function or reduce the risk of infection (Kaminogawa & Nanno, 2004; Goldsmith & Sartor, 2014). Studies have suggested that diet can improve depressed immune function by moderating the severity of infectious diseases and reducing infection rates (El-Gamal et al., 2011; Rajilić-

Stojanović et al., 2015). Cereals are staple foods and feed more than half of the world's population; they are composed mainly of starch, protein, some minerals, and non-starch polysaccharides that cannot be digested by human enzymes (Mohan et al., 2010). Arabinoxylan is the main non-starch polysaccharide of many cereals (Zhou et al., 2010; Lovegrove et al., 2017). It has been reported that arabinoxylan possesses immune-modulatory effects (Li et al., 2015). Moreover, there is a structural similarity between arabinoxylans and the lipopolysaccharides of Gram-negative bacteria (Park et al., 2009; Park et al., 2017). The aim of this review is to give an overview of the immune system and how arabinoxylans might modulate the immune response.

2. Immune system

The immune system of the human body is a complex network of molecules, cells and organs that interact and communicate together to respond to the invasion of pathogens and maintain the body's homoeostasis (Thompson, 2015). The immune system consists of innate immunity, which is a stereotyped rapid response to a stimulus, and adaptive immunity, which is a slower but highly specific response (Iwasaki & Medzhitov, 2015). The innate immune response operates in conjunction with the adaptive through activation of signalling pathways (O'Neill et al., 2013). Figure 1 gives an overview of the interactions between the innate and adaptive responses of the immune system.

2.1. Innate immune system

The innate immune system is the first line of defence in the human body and is composed of three stages (Orlowsky & Kraus, 2015). The first stage is the chemical and physical barriers while the second stage depends on cell-intrinsic mechanisms and digestive enzymes which destroy viruses, bacteria and other pathogenic invaders (Medzhitov & Janeway Jr, 2000; Mogensen, 2009; Iwasaki & Pillai, 2014; Iwasaki & Medzhitov, 2015). The third stage of defence relies on the recognition of preserved pathogenic features (pathogen-associated immune-stimulants) by the complementary system and phagocytosis by immune cells such as natural killer cells, neutrophils and macrophages (McCarthy et al., 2014). Pathogenic immune stimulants are referred to as Pathogen-Associated Molecular Patterns (PAMPs) and they include pathogen cell wall polysaccharides such as chitin and mannans from fungi, lipopolysaccharides (LPS) from Gram-negative bacteria and peptidoglycan from Gram-positive bacteria (Volman et al., 2008; Kawai & Akira, 2010; Wiersinga et al., 2014).

PAMPs have been well studied, especially LPS (Okuda et al., 2016). This has the ability to initiate the host defence through recognition of its bioactive component, lipid A, via co-receptor

MD-2 and Toll-Like Receptor 4 (TLR4) (Saitoh et al., 2004; Ohto et al., 2012; Kang et al., 2016). The structure of lipid A is composed of two glucosamine units with a $\beta(1\rightarrow6)$ linkage attached to six fatty acyl chains, 1 and 4 phosphate groups (Slocum et al., 2014; Trouw et al., 2017). It has been reported that optimal immune activation of lipid A is derived from the acyl chains attached directly to the di-glucosamine (Raetz et al., 2009).

The complement system is part of the innate immune system (Galluzzi et al., 2017) and is responsible for enhancing the ability of phagocytic cells to clear damaged cells and microbes from the system (Orsini et al., 2014). Three pathways have been identified for complement activation, which are the classical, alternative and lectin pathway (Takahashi et al., 2008; Merle et al., 2015). IgG or IgM antigen/antibody complexes are responsible for initiating the classical pathway through binding to the first protein of the cascade (C1q) which in turn activates the C1r, leading to formation of the membrane attack complex which eventually penetrates bacterial membranes creating pores which lead to bacterial lysis (Peerschke et al., 2016).

The second pathway of the complement system is known as the alternative pathway or ‘properdin pathway’, which is a failure to regulate low-level continuous formation of C3 convertase (Miwa et al., 2013). Eventually, if a product of C3 convertase called C3b binds to a bacterial cell surface, this creates an amplification loop for other pathways (Galluzzi et al., 2017; Trouw et al., 2017).

The lectin pathway involves mannose-binding lectin (MBL). The initiating molecules for this pathway (MBL and ficolin) are multimeric lectin complexes (Amiri, 2015). These molecules bind to specific carbohydrates in the host to activate the pathway through enzymatic activity of mannan-binding lectin-associated serine protease (MASP). The structural similarities between C1 and MBL suggest that complement activation by C1 and MBL involves similar pathways (Kozarcanin et al., 2016).

All three pathways have the ability to activate the key components C 1-3, referred to as C3. This activation is critical for the complementary reaction as it triggers the inflammatory response, which in turn activates components C 5-9 (Ali et al., 2012).

Activation of C 5-9 triggers a cascade of events that leads to activation and recruitment of other innate immune cells (Garred et al., 2016). The PAMPs from the invaders bind to Pattern Recognition Receptors (PRRs) which are displayed on host immune cells (Takeuchi & Akira, 2010; Kagan & Barton, 2015). PRRs include the Toll-Like Receptors (TLRs) which are found on the surface of phagocytes (dendritic cells, neutrophils and macrophages) (De Nardo, 2015). For example, Toll-Like Receptor-4 (TLR4) activates the innate immune response through recognition of LPS from the cell wall of Gram-negative bacteria (He et al., 2014).

Post-activation, immune cells such as neutrophils, dendritic cells and macrophages secrete cytokines to communicate with other cells in the immune system and stimulate the immune response (Guilliams et al., 2014). On the other hand, activation of innate immune cells produces digestive enzymes and reactive oxygen radicals that destroy pathogens (Takahashi et al., 2008). Furthermore, dendritic cells play an important role in transferring ingested pathogens to the lymph nodes to activate T lymphocytes, thereby initiating a specific immune response that is part of the adaptive immune system (Kim et al., 2006; Ait-Oufella et al., 2014).

2.2. Adaptive immune system

Adaptive immunity protects the human body from certain death by infection (Wong et al., 2014). Once the innate response is initiated, it calls the adaptive immune system into play; then both work together to eliminate pathogens (Zhu et al., 2015). Unlike the innate immune response, the adaptive immune response is slow but highly specific against pathogens, and its protection is long-lasting (Cooper & Alder, 2006). It is pointless to mount the adaptive immune response against harmless foreign molecules, otherwise the adaptive immune response might

be deleterious (Witztum & Lichtman, 2014). This is normally avoided because the adaptive immune response is triggered by the innate immune system only when the latter recognises molecules of the attacking pathogens such as PAMPs (Bonneaud et al., 2003; Lu et al., 2016). Dendritic cells, also known as pathogen-presenting cells, have surfaces packed with PRRs, which bind to the PAMPs of foreign pathogens and initiate phagocytosis (Visintin et al., 2001; Gringhuis et al., 2014). The dendritic cells, with the ingested pathogens, then move to a peripheral lymphoid organ or to a nearby lymph node where the dendritic cells present the antigens of the ingested pathogens to T lymphocytes (Cravens & Lipsky, 2002; Gringhuis et al., 2014). The cell surface of T cells is covered with various receptors that recognise extraneous antigens such as foreign polysaccharides or large proteins (Caramalho et al., 2003; Levine, 2015). To complete the activation of T cells, a co-stimulatory signal is sent from the dendritic cells, resulting in proliferation of T cells with the same receptor, thereby inducing antigen-specific adaptive immune responses (Chen & Flies, 2013). To eliminate the pathogen at the infection site, T cells mature and differentiate into different types of effector T cells including cytotoxic, helper and regulatory T cells (Mucida et al., 2013; Nishikawa & Sakaguchi, 2014). Cytotoxic T cells have the ability to detect substantial number of different antigens with high specificity and release lytic proteins thus eliminating pathogens that proliferate inside the host cell (Jones et al., 2017).

Helper T cells can release cytokines that guide dendritic cells to stay in their active form and can stimulate antibody production from B cells that kill pathogens (Rissoan et al., 1999; Ise et al., 2014). Helper T cells can also express co-stimulatory proteins on their surface and release cytokines to activate more cytotoxic T cells and macrophages (Croft, 2003; Maude et al., 2014). The regulation and control of activated immune cells is mediated by regulatory T cells, which can inhibit the activity of cytotoxic T cells, helper T cells and dendritic cells to avoid autoimmunity (Sakaguchi et al., 2008; Ito et al., 2016). Activated B cells secrete serum proteins

and synthesize antibodies that bind directly to pathogens to inactivate them. They also recruit innate immune cells such as macrophages to eliminate invaders (Clark & Ledbetter, 1994; Amable et al., 2014).

2.3. Monocytes

Monocytes are white cells circulating in the blood (Guilliams et al., 2014). They can express CD11b and Toll-like receptor-4 (TLR4) associated with CD14, which are triggered by LPS from the cell wall of Gram-negative bacteria (Taylor et al., 2005; Frantz et al., 2013). Monocytes originate from haematopoietic stem cells in the bone marrow (Lee et al., 2015) and activation of these stem cells results in the differentiation of common myeloid progenitors (CMPs) which differentiate further into macrophage and granulocyte progenitors (Ogawa, 1993; Ginhoux & Jung, 2014). Prior to the transformation of haematopoietic stem cells into circulating monocytes, they undertake a series of embryonic divisions (Chow et al., 2011; Lim et al., 2013). Monocytes circulate in the blood and have the potential to differentiate into dendritic cells or macrophages (Geissmann et al., 2010; Heidt et al., 2014). Differentiation to macrophages requires the activation of runt-related transcription factor that encodes the ETS family transcription factor PU.1, which needs to be constantly expressed at high levels to induce monocyte differentiation to macrophages (Lawrence & Natoli, 2011; Schneider et al., 2014).

2.4. Macrophages

Monocytes circulate in the blood for up to two days followed by migration into tissues and differentiation into macrophages (van Furth & Cohn, 1968; Geissmann et al., 2010). Macrophages and neutrophils have the ability to take up pathogens through phagocytosis, a process which engulfs large particles ($> 0.5 \mu\text{m}$) into cells through an actin-dependent

mechanism (Silva, 2010; Linehan et al., 2014). Activation of macrophages and subsequent phagocytosis occurs through PAMP-mediated recognition of Gram-negative and Gram-positive bacteria by PRRs (Plüddemann et al., 2011; Martinez & Gordon, 2014). Macrophages have a range of PRRs including TLR, which when activated results in pro-inflammatory cytokine production including IL-23, IL-12, IL-6 and TNF α (Mosser & Edwards, 2008; O'Neill & Pearce, 2016). Activated macrophages also express inducible nitric oxide synthase (iNOS), which is responsible for generating nitric oxide (NO), a key mediator for killing bacteria within macrophages (Murray et al., 2014; Martins et al., 2017).

2.5. Initiate immune responses

PAMPs are responsible for initiating the innate immune response through PRRs, of which the TLR family has been extensively investigated in recent years (Medzhitov, 2001; Akira & Takeda, 2004; Pradere et al., 2014). On recognition of PAMPs, PRRs at the cell surface trigger intracellular pathways that lead to the transcription of chemokines and cytokines involved in antimicrobial and proinflammatory responses (Akira & Takeda, 2004; Gazendam et al., 2016).

2.6. Toll-like receptor family (TLRs)

The most investigated class of PRRs is the TLR family. The name is derived from their homology to the Toll protein in *Drosophila melanogaster* (Medzhitov et al., 1997; De Nardo, 2015; Murofushi et al., 2015). The structural hallmarks of all known TLRs are an extracellular cysteine-rich domain, leucine-rich motifs (LRR) and a cytoplasmic signalling Toll/IL-1 receptor (TIR) analogy domain in the intracellular region (Gay & Keith, 1991; Pietretti et al., 2014; Ren et al., 2014). The intracellular signal transduction is due to receptor oligomerisation that is induced by ligand binding to TLRs (Futosi et al., 2013). To date, 10 TLRs have been identified in mammals, and each recognize distinct PAMPs derived from bacteria, viruses, fungi and protozoa (Akira & Takeda, 2004; Bonham et al., 2014). The TLRs include TLR1,

TLR2, TLR4 and TLR6 that recognize lipoproteins such as triacyl lipopeptides and LPS, while TLR3, TLR7, TLR8 and TLR9 recognize nucleic acids such as dsRNA or ssRNA (Curtale et al., 2013; Vacchelli et al., 2013; Gay et al., 2014).

2.7. Lipopolysaccharides (LPS)

LPS are a major component of the cell wall of Gram-negative bacteria (Guha & Mackman, 2001), consisting of three parts; a polysaccharide side chain also known as O-antigen or O-chain, a non-repeating core polysaccharide, and lipid A, which is hydrophobic (Speciale et al., 2015). The polysaccharide side chain and non-repeating core polysaccharide are projections from the surface, while the hydrophobic domain is embedded in the outer membrane. The lipid A domain is a source of toxicity, while the O-chains are easily detected by the host antibodies and, to avoid detection, are often modified by bacteria (Lerouge & Vanderleyden, 2002; Miller et al., 2005; Eckert et al., 2013; Wu et al., 2013). Lipopolysaccharide structure is illustrated in Figure 2. Low levels of LPS are sufficient to induce a substantial inflammatory response of the innate immune system (Schwarz et al., 2014). LPS binds to the LPS Binding Protein (LBP) in serum, before being transferred to CD14 and then to MD2, which is associated with TLR4 (Ryu et al., 2017). The receptor complex then promotes secretion of nitric oxide (NO) and pro-inflammatory cytokines such as TNF α and IL-8 in monocytes and macrophages (Johnson et al., 2002; Termeer et al., 2002; Miller et al., 2005; Massey et al., 2015; Lee et al., 2016).

2.8. Cytokines

Cytokines are small, soluble proteins that affect the function or growth of cells. Cytokines can act in a paracrine way (affect nearby cells) or an autocrine way (affect the same cell). However, some cytokines such as IL-6, IL-8 and TNF α can have systemic effects. Cytokines act on immune cells and mediate inflammatory responses (Vilček & Feldmann, 2004; Le Maitre, 2014).

2.9. Tumour necrosis factor alpha (TNF α)

TNF α is a pro-inflammatory cytokine with various biological effects (Bekkering et al., 2014). TNF α is involved in apoptosis, differentiation and proliferation (Du et al., 2014; Sullivan et al., 2014) and levels are elevated in inflammatory diseases such as rheumatoid arthritis (Motley et al., 2004; Garraway et al., 2014). Two receptors are known to mediate the effects of TNF α , TNFR1 and TNFR2. Local TNF α production is critical for elimination of local infections (Motley et al., 2004; Olmos & Lladó, 2014). Systemic TNF α release also plays a vital role in septic shock (Kanashiro et al., 2017). TNF α is released in macrophages and monocytes in response to foreign stimuli such as LPS from Gram-negative bacteria. The secretion of TNF α from T-cells is initiated by activation of the T-cell receptor (Manzo et al., 2017). In addition, natural killer cells and B cells can produce TNF α (Eissner et al., 2000; Yu et al., 2009). The effect of TNF α on endothelial cells includes the upregulation of leukocyte adhesion molecules that contribute to leukocyte recruitment (Huang et al., 2015).

2.10. Nitric oxide (NO)

Nitric oxide (NO) is a short-lived, gaseous, small molecule composed of one atom of oxygen and one atom of nitrogen, thus making it a free radical due to unpaired electrons (Pacher et al., 2007). In the human body, NO is defined as a product of macrophage activation by pro-inflammatory cytokines, microbial endotoxins such as LPS, or both (Rath et al., 2014). NO is a product of L-arginine degradation, the reaction being catalysed by an enzyme called inducible nitric oxide synthase (iNOS) (Bogdan, 2001; Palygin et al., 2015). This reaction requires several cofactors including calcium/calmodulin, flavin mononucleotide (FMN), nicotinamide adenine dinucleotide phosphate (NADPH), flavin adenine dinucleotide (FAD), tetrahydrobiopterin (BH₄), oxygen and haem (Baek et al., 1993; Marletta, 1994). There are three known isoforms of NOS; their names come from where they were first found. Inducible

NOS (iNOS) was found in macrophages, endothelial NOS (eNOS) found in endothelial cells and the neuronal NOS (nNOS) found in the brain. These isoforms are also known as NOS-2, NOS-3 and NOS-1 respectively (Alderton et al., 2001; Bogdan, 2015).

2.10.1. NO role in macrophages

NO production in macrophages depends on their activation in response to cytokine or bacterial endotoxin stimuli (Sadek et al., 2017). Macrophages act as patrolling cells and produce low levels of NO in quiescent conditions (Prolo et al., 2015). However, once activated by stimuli, macrophages produce excessive amounts of NO, releasing NO₂⁻ and NO₃⁻, which are important in pathogen scavengers (Marletta et al., 1988; Hibbs, 2002). Although NO production is critical for macrophage phagocytosis, an excessive amount of NO has been associated with high levels of necrosis and apoptosis (He et al., 2016; Jakubowska et al., 2016). It has been reported that high levels of NO induce autoimmune reactions such as asthma and arthritis (Scichilone et al., 2013), so it is crucial to dampen NO production in such conditions. Evidence suggests the modulation of immune function can be achieved by food (Zeng et al., 2016).

2.11. Immunomodulating properties of food

The functionality of the immune system is crucial for protecting the body from the attacks of pathogens or from cancer cell proliferation (Mittal et al., 2014). Thus, it plays a vital role in health homeostasis (Nairz et al., 2013). However, many factors disturb immune functions such as an unhealthy lifestyle, malnutrition, physical stress, disease and ageing (Nahrendorf & Swirski, 2015). Evidence suggests that the ingestion of foods with immuneomodulatory effects prevents declining immune function or reduces the risk of infection (Kaminogawa & Nanno, 2004; Goldsmith & Sartor, 2014). There are many studies suggesting that foods can improve depressed immune function by moderating the severity of infectious diseases and reducing infection rates (El-Gamal et al., 2011; Rajilić-Stojanović et al., 2015). Thus, foods with the

ability to improve immune responses, particularly in patients with impaired immunity such as cancer (Braga et al., 1996; Bradbury et al., 2014; Schnekenburger et al., 2014), are both clinically and commercially valuable.

3. Arabinoxylans

Arabinoxylan is a non-starch polysaccharide (Figure 3) with a backbone of β -(1-4)-linked d-xylopyranosyl residues to which α -l-arabinofuranose units are linked as side chains in the second and/or third carbon-positions (Courtin et al., 2000; Roubroeks et al., 2000; Zhou et al., 2010; Qiu et al., 2017).

Arabinoxylan structure is characterised by substitution of the xylopyranose- linked xylan backbone. L-Arabinofuranose is the main sugar substitute for xylopyranose residues at O-2 and/or O-3 via α -1, 2 and α - 1, 3 glycosidic linkages. This leads to three different forms namely, un-substituted xylopyranose, mono-substituted xylopyranose at O-2 or O-3 and di-substituted xylopyranose at O-2 and O-3 (Izydorczyk & Biliaderis, 1995; Saulnier et al., 2007; Broekaert et al., 2011; Qiu et al., 2017). However, arabinofuranose substitutions can form short oligosaccharide side chains that comprise of two or more arabinofuranose residues (Figure 4) (Saulnier et al., 2007).

There are many techniques available for arabinoxylan extraction and different extraction methods give different yields and degrees of branching, molecular weight distribution and tertiary conformation (Lu et al., 2005), i.e. hot water extraction (Izydorczyk et al., 1998; Cyran et al., 2003; Iqbal et al., 2011; Yu et al., 2017) and ultrasound-assisted enzymatic extraction (Wang et al., 2014).

Arabinoxylans can be classified according to their solubility in water, as either water-unextractable arabinoxylans (WUAX) or water-extractable arabinoxylans (WEAX) (Moers et al., 2005). The structure of WUAX is different from that of WEAX as WUAX will not

solubilise in water, but will be solubilized in alkaline solutions (Gruppen et al., 1993). Table 1 shows the water-extractable and water-unextractable arabinoxylans in some cereals.

Reducing the molecular weight of the arabinoxylans not only increases their solubility in water but also increases their biological health benefits (Li et al., 2013). Recently, pronounced effects of low Mw arabinoxylans (66 kDa) have been observed to have a higher prebiotic stimulation in an *in vitro* study compared to higher Mw arabinoxylans (Hughes et al., 2007). Modification of the molecular characteristics of arabinoxylans such as Mw is important to achieve the optimum prebiotic, anti-tumour activities and immune stimulation (Li et al., 2013).

It has been suggested that the activity of arabinoxylans is dependent on their sugar composition, molecular weight and degree of branching (Zhou et al., 2010; Cao et al., 2011). The most investigated type of arabinoxylans (MGN-3) has low molecular weight with a low arabinose-to-xylose ratio (0.5) (Zhang et al., 2015) which is similar to the enzyme- extracted wheat bran arabinoxylans (Zhou et al., 2010). Both polysaccharides could activate macrophages (Zhou et al., 2010; Zhang et al., 2015). However, MGN-3 appeared to be more effective, which might be due to differences in the sugar composition since MGN-3 has more glucose and galactose side chains (Zhang et al., 2015). Figure 5. shows a simplified representation of the macrophage TLR 4 receptor with arabinoxylans and LPS.

3.1. Immunomodulatory potentials of arabinoxylans

It has been reported that oral administration of arabinoxylans extracted from wheat bran using xylanases and alkali extraction has an immune-modulatory effect on both the innate and adaptive immune systems (Zhou et al., 2010; Cao et al., 2011). Alkali extracted arabinoxylans from wheat bran showed inhibitory effects on tumour growth and IL-2 production at 100-400 mg/kg on S 180 tumour bearing mice. The most significant results were at the highest

concentration (400 mg/kg). Moreover, there was an increase in leukocyte count, and stem cell proliferation was enhanced after oral administration (Cao et al., 2011).

Another study conducted by Zhou et al. (2010) indicated that (200 mg/kg) oral administration of enzyme-extracted wheat bran arabinoxylans exhibited immunostimulatory effects on both innate and adaptive immunity. The enzyme-extracted arabinoxylans stimulated phagocytosis by macrophages and delayed hypersensitivity more than alkaline-extracted arabinoxylans (Zhou et al., 2010).

Recently, Li et al. (2015) investigated the effect of enzyme-extracted arabinoxylans (AXE) from wheat endosperm pentosan on U937 and Caco-2 cell lines. They reported that AXE generated higher nitric oxide (NO) levels than (WEAX) and the increase in NO production was dose-dependent. AXE was reported to be more effective than WEAX in stimulating IL-8 production (Li et al., 2015).

Previous studies suggested that arabinoxylans from various sources have immunomodulatory potential. It is clear that there are several factors affecting the immunomodulatory potential of arabinoxylans including the method of extraction, enzyme/chemical treatments and botanical source. Table 2 shows the structural activity of arabinoxylans from rice bran and wheat on different cells.

3.2. Potential arabinoxylan receptors

Although receptors for arabinoxylans have not been identified, some potential receptors have been proposed. AXs maybe acting like PAMPs since AXs from cornhusk and rice bran have shown similarities in terms of molecular weight and structure to LPS from Gram-negative bacteria (Ghoneum & Brown, 1999; Ogawa et al., 2005; Mendis et al., 2017). For example, LPS has outer core hexoses such as glucose and galactose, which are found in AXs. Another similarity between the structure of LPS and AXs includes C-3 branched polysaccharides

(Rietschel et al., 1994; Heinrichs et al., 1998). Moreover, AXE from wheat endosperm pentosan had low Mw (1-25 kDa) (Li et al., 2015) which is within the molecular weight range (13-20 kDa) of LPS (Mangoni et al., 2008). Therefore, arabinoxylans may activate phagocytes by attaching to Toll-like receptors expressed on their cell surface (Zhang et al., 2015). Since LPS specifically binds to TLR4, it suggests that TLR4 may also be a potential receptor for arabinoxylans (Mendis et al., 2016). If true, arabinoxylans may compete with LPS for the TLR4 receptor during infection, thus mediating the LPS-induced immune response.

Other receptors besides TLRs may also act as receptors for arabinoxylans. These include the Dectin-1 receptor, which has been reported to be a β glucan receptor (Karumuthil-Melethil et al., 2014; Kanjan et al., 2017). However, Sahasrabudhe et al. (2016) have reported recently that arabinoxylan from wheat has the ability to stimulate Dectin-1 receptors and it enhanced Interleukin 23 (IL-23), and Interleukin 4 (IL-4) expression in Dectin-1 stimulated dendritic cells (Sahasrabudhe et al., 2016).

3.3. Structure-activity relationship

The most investigated type of arabinoxylans (MGN-3) has a low molecular weight with a low arabinose- to-xylose ratio (0.5) (Zhang et al., 2015) which is similar to the enzyme- extracted wheat bran arabinoxylans (Zhou et al., 2010). Both polysaccharides appear to activate macrophages but MGN-3 appeared to be more effective, which might be due to the higher sugar composition (Zhang et al., 2015). Figure 5. shows a simplified representation of the macrophage TLR 4 receptor with arabinoxylans and LPS.

4. Conclusions

Cereals are by far the most important source of food all over the world. Arabinoxylans are the major non-starch polysaccharides in most of the cereals. Also, it has been reported that

396 arabinoxylans have immune-modulatory activities. The immunomodulatory potentials of
397 arabinoxylans have been linked to their sugar composition, molecular weight and degree of
398 branching. Furthermore, there are structural similarities between arabinoxylans and LPS in
399 terms of molecular weight and structure, suggesting that arabinoxylans can modulate the
400 immune response through activation the LPS receptor TLR 4. Future work should focus on
401 understanding more of the arabinoxylan mechanism of action, which might help in modulating
402 the immune response more efficiently.

References

- Ait-Oufella, H., Sage, A. P., Mallat, Z., & Tedgui, A. (2014). Adaptive (T and B cells) immunity and control by dendritic cells in atherosclerosis. *Circulation research*, 114(10), 1640-1660.
- Akira, S., & Takeda, K. (2004). Toll-like receptor signalling. *Nature Reviews Immunology*, 4(7), 499-511.
- Alderton, W. K., Cooper, C. E., & Knowles, R. G. (2001). Nitric oxide synthases: structure, function and inhibition. *Biochemical Journal*, 357(3), 593-615.
- Ali, Y. M., Lynch, N. J., Haleem, K. S., Fujita, T., Endo, Y., Hansen, S., Holmskov, U., Takahashi, K., Stahl, G. L., & Dudler, T. (2012). The lectin pathway of complement activation is a critical component of the innate immune response to pneumococcal infection. *PLoS Pathog*, 8(7), e1002793.
- Amable, P. R., Teixeira, M. V. T., Carias, R. B. V., Granjeiro, J. M., & Borojevic, R. (2014). Protein synthesis and secretion in human mesenchymal cells derived from bone marrow, adipose tissue and Wharton's jelly. *Stem cell research & therapy*, 5(2), 53-66.
- Amiri, F. S. (2015). Role of ficolin-3 in acute kidney graft rejection: A new diagnostic tool. *Indian Journal of Transplantation*, 9(4), 164-167.
- Baek, K. J., Thiel, B. A., Lucas, S., & Stuehr, D. J. (1993). Macrophage nitric oxide synthase subunits. Purification, characterization, and role of prosthetic groups and substrate in regulating their association into a dimeric enzyme. *Journal of Biological Chemistry*, 268(28), 21120-21129.
- Bekkering, S., Quintin, J., Joosten, L. A., van der Meer, J. W., Netea, M. G., & Riksen, N. P. (2014). Oxidized low-density lipoprotein induces long-term proinflammatory cytokine production and foam cell formation via epigenetic reprogramming of monocytes. *Arteriosclerosis, thrombosis, and vascular biology*, ATVB AHA. 114.303887.
- Bogdan, C. (2001). Nitric oxide and the immune response. *Nature immunology*, 2(10), 907-916.
- Bogdan, C. (2015). Nitric oxide synthase in innate and adaptive immunity: an update. *Trends in immunology*, 36(3), 161-178.
- Bonham, K. S., Orzalli, M. H., Hayashi, K., Wolf, A. I., Glanemann, C., Weninger, W., Iwasaki, A., Knipe, D. M., & Kagan, J. C. (2014). A promiscuous lipid-binding protein diversifies the subcellular sites of toll-like receptor signal transduction. *Cell*, 156(4), 705-716.
- Bonneaud, C., Mazuc, J., Gonzalez, G., Haussy, C., Chastel, O., Faivre, B., & Sorci, G. (2003). Assessing the cost of mounting an immune response. *The American Naturalist*, 161(3), 367-379.
- Bradbury, K., Balkwill, A., Spencer, E., Roddam, A. W., Reeves, G. K., Green, J., Key, T., Beral, V., Pirie, K., & Banks, E. (2014). Organic food consumption and the incidence of cancer in a large prospective study of women in the United Kingdom. *British journal of cancer*, 110(9), 2321-2326.
- Braga, M., Vignali, A., Gianotti, L., Cestari, A., Profili, M., & Carlo, V. (1996). Immune and nutritional effects of early enteral nutrition after major abdominal operations. *The European journal of surgery= Acta chirurgica*, 162(2), 105-112.
- Broekaert, W. F., Courtin, C. M., Verbeke, K., Van de Wiele, T., Verstraete, W., & Delcour, J. A. (2011). Prebiotic and other health-related effects of cereal-derived arabinoxylans, arabinoxylan-oligosaccharides, and xylooligosaccharides. *Critical Reviews in Food Science and Nutrition*, 51(2), 178-194.

- Cao, L., Liu, X., Qian, T., Sun, G., Guo, Y., Chang, F., Zhou, S., & Sun, X. (2011). Antitumor and immunomodulatory activity of arabinoxylans: A major constituent of wheat bran. *International Journal of Biological Macromolecules*, 48(1), 160-164.
- Caramalho, I., Lopes-Carvalho, T., Ostler, D., Zelenay, S., Haury, M., & Demengeot, J. (2003). Regulatory T cells selectively express toll-like receptors and are activated by lipopolysaccharide. *The Journal of experimental medicine*, 197(4), 403-411.
- Chen, L., & Flies, D. B. (2013). Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nature Reviews Immunology*, 13(4), 227-242.
- Choct, M. (1997). Feed non-starch polysaccharides: chemical structures and nutritional significance. *Feed milling international*, 191(June issue), 13-26.
- Chow, A., Brown, B. D., & Merad, M. (2011). Studying the mononuclear phagocyte system in the molecular age. *Nature Reviews Immunology*, 11(11), 788-798.
- Clark, E. A., & Ledbetter, J. A. (1994). How B and T cells talk to each other. *Nature*, 367(6462), 425-428.
- Cooper, M. D., & Alder, M. N. (2006). The evolution of adaptive immune systems. *Cell*, 124(4), 815-822.
- Courtin, C., Van den Broeck, H., & Delcour, J. (2000). Determination of reducing end sugar residues in oligo- and polysaccharides by gas-liquid chromatography. *Journal of Chromatography A*, 866(1), 97-104.
- Cravens, P. D., & Lipsky, P. E. (2002). Dendritic cells, chemokine receptors and autoimmune inflammatory diseases. *Immunology and cell biology*, 80(5), 497-505.
- Croft, M. (2003). Co-stimulatory members of the TNFR family: keys to effective T-cell immunity? *Nature Reviews Immunology*, 3(8), 609-620.
- Curtale, G., Mirolo, M., Renzi, T. A., Rossato, M., Bazzoni, F., & Locati, M. (2013). Negative regulation of Toll-like receptor 4 signaling by IL-10-dependent microRNA-146b. *Proceedings of the National Academy of Sciences*, 110(28), 11499-11504.
- Cyran, M., Courtin, C. M., & Delcour, J. A. (2003). Structural features of arabinoxylans extracted with water at different temperatures from two rye flours of diverse breadmaking quality. *Journal of agricultural and food chemistry*, 51(15), 4404-4416.
- De Nardo, D. (2015). Toll-like receptors: Activation, signalling and transcriptional modulation. *Cytokine*, 74(2), 181-189.
- Dodevska, M. S., Djordjevic, B. I., Sobajic, S. S., Miletic, I. D., Djordjevic, P. B., & Dimitrijevic-Sreckovic, V. S. (2013). Characterisation of dietary fibre components in cereals and legumes used in Serbian diet. *Food Chemistry*, 141(3), 1624-1629.
- Dranoff, G. (2004). Cytokines in cancer pathogenesis and cancer therapy. *Nature Reviews Cancer*, 4(1), 11-22.
- Du, G., Song, Y., Zhang, T., Ma, L., Bian, N., Chen, X., Feng, J., Chang, Q., & Li, Z. (2014). Simvastatin attenuates TNF- α -induced apoptosis in endothelial progenitor cells via the upregulation of SIRT1. *International journal of molecular medicine*, 34(1), 177-182.
- Eckert, J. K., Kim, Y. J., Kim, J. I., Gürtler, K., Oh, D.-Y., Sur, S., Lundvall, L., Hamann, L., van der Ploeg, A., & Pickkers, P. (2013). The crystal structure of lipopolysaccharide binding protein reveals the location of a frequent mutation that impairs innate immunity. *Immunity*, 39(4), 647-660.
- Edwards, S., Chaplin, M. F., Blackwood, A. D., & Dettmar, P. W. (2003). Primary structure of arabinoxylans of ispaghula husk and wheat bran. *Proceedings of the Nutrition Society*, 62(01), 217-222.
- Eissner, G., Kirchner, S., Lindner, H., Kolch, W., Janosch, P., Grell, M., Scheurich, P., Andreesen, R., & Holler, E. (2000). Reverse Signaling Through Transmembrane TNF Confers Resistance to Lipopolysaccharide in Human Monocytes and Macrophages. *The Journal of Immunology*, 164(12), 6193-6198. doi: 10.4049/jimmunol.164.12.6193

- El-Gamal, Y. M., Elmasry, O. A., El-Ghoneimy, D. H., & Soliman, I. M. (2011). Immunomodulatory effects of food. *Egyptian Journal of Pediatric Allergy and Immunology (The)*, 9(1).
- Frantz, S., Hofmann, U., Fraccarollo, D., Schäfer, A., Kranepuhl, S., Hagedorn, I., Nieswandt, B., Nahrendorf, M., Wagner, H., & Bayer, B. (2013). Monocytes/macrophages prevent healing defects and left ventricular thrombus formation after myocardial infarction. *The FASEB Journal*, 27(3), 871-881.
- Futosi, K., Fodor, S., & Mócsai, A. (2013). Reprint of Neutrophil cell surface receptors and their intracellular signal transduction pathways. *International immunopharmacology*, 17(4), 1185-1197.
- Galluzzi, L., Buqué, A., Kepp, O., Zitvogel, L., & Kroemer, G. (2017). Reply: The complement system is also important in immunogenic cell death. *Nature Reviews Immunology*, 17(2), 143-151.
- Garófalo, L., Vazquez, D., Ferreira, F., & Soule, S. (2011). Wheat flour non-starch polysaccharides and their effect on dough rheological properties. *Industrial Crops and Products*, 34(2), 1327-1331.
- Garraway, S. M., Woller, S. A., Huie, J. R., Hartman, J. J., Hook, M. A., Miranda, R. C., Huang, Y.-J., Ferguson, A. R., & Grau, J. W. (2014). Peripheral noxious stimulation reduces withdrawal threshold to mechanical stimuli after spinal cord injury: role of tumor necrosis factor alpha and apoptosis. *PAIN®*, 155(11), 2344-2359.
- Garred, P., Genster, N., Pilely, K., Bayarri-Olmos, R., Rosbjerg, A., Ma, Y. J., & Skjoedt, M. O. (2016). A journey through the lectin pathway of complement—MBL and beyond. *Immunological reviews*, 274(1), 74-97.
- Gay, N. J., & Keith, F. J. (1991). Drosophila Toll and IL-1 receptor. *Nature*, 351, 355-356.
- Gay, N. J., Symmons, M. F., Gangloff, M., & Bryant, C. E. (2014). Assembly and localization of Toll-like receptor signalling complexes. *Nature Reviews Immunology*, 14(8), 546-558.
- Gazendam, R. P., van de Geer, A., van Hamme, J. L., Helgers, L., Rohr, J., Chrabieh, M., Picard, C., Roos, D., van den Berg, J. M., & van den Berg, T. (2016). Proinflammatory cytokine response toward fungi but not bacteria in chronic granulomatous disease. *Journal of Allergy and Clinical Immunology*, 138(3), 928-930. e924.
- Geissmann, F., Manz, M. G., Jung, S., Sieweke, M. H., Merad, M., & Ley, K. (2010). Development of monocytes, macrophages, and dendritic cells. *Science*, 327(5966), 656-661.
- Ghoneum, M., & Brown, J. (1999). NK immunorestitution of cancer patients by biobran/MGN-3, a modified arabinoxylan rice bran (study of 32 patients followed for up to 4 year). *Anti Aging Med. Ther*, 3, 217-226.
- Ginhoux, F., & Jung, S. (2014). Monocytes and macrophages: developmental pathways and tissue homeostasis. *Nature Reviews Immunology*, 14(6), 392-404.
- Goldsmith, J. R., & Sartor, R. B. (2014). The role of diet on intestinal microbiota metabolism: downstream impacts on host immune function and health, and therapeutic implications. *Journal of gastroenterology*, 49(5), 785-798.
- Gringhuis, S. I., Kaptein, T. M., Wevers, B. A., Van Der Vlist, M., Klaver, E. J., Van Die, I., Vriend, L. E., De Jong, M. A., & Geijtenbeek, T. B. (2014). Fucose-based PAMPs prime dendritic cells for follicular T helper cell polarization via DC-SIGN-dependent IL-27 production. *Nature communications*, 5.
- Gruppen, H., Kormelink, F., & Voragen, A. (1993). Water-unextractable cell wall material from wheat flour. 3. A structural model for arabinoxylans. *Journal of Cereal Science*, 18(2), 111-128.

- Guha, M., & Mackman, N. (2001). LPS induction of gene expression in human monocytes. *Cellular signalling*, 13(2), 85-94.
- Guilliams, M., Ginhoux, F., Jakubzick, C., Naik, S. H., Onai, N., Schraml, B. U., Segura, E., Tussiwand, R., & Yona, S. (2014). Dendritic cells, monocytes and macrophages: a unified nomenclature based on ontogeny. *Nature Reviews Immunology*, 14(8), 571-578.
- Hansen, H. B., Rasmussen, C. V., Bach Knudsen, K. E., & Hansen, Å. (2003). Effects of genotype and harvest year on content and composition of dietary fibre in rye (*Secale cereale* L) grain. *Journal of the Science of Food and Agriculture*, 83(1), 76-85.
- Hashimoto, S., Shogren, M., & Pomeranz, Y. (1987). Cereal pentosans: their estimation and significance. I. Pentosans in wheat and milled wheat products. *Cereal Chemistry*, 64(1), 30-34.
- Hashimoto, S., Shogren, M., Bolte, L., & Pomeranz, Y. (1987). Cereal pentosans: their estimation and significance. 3. Pentosans in abraded grains and milling by-products. *Cereal Chemistry*, 64(1), 39-41.
- He, B., Tao, H., Wei, A., Liu, S., Li, X., & Chen, R. (2016). Protection of carboxymethylated chitosan on chondrocytes from nitric oxide-induced apoptosis by regulating phosphatidylinositol 3-kinase/Akt signaling pathway. *Biochemical and biophysical research communications*, 479(2), 380-386.
- He, W., Wang, Z., Zhou, Z., Zhang, Y., Zhu, Q., Wei, K., Lin, Y., Cooper, P. R., Smith, A. J., & Yu, Q. (2014). Lipopolysaccharide enhances Wnt5a expression through toll-like receptor 4, myeloid differentiating factor 88, phosphatidylinositol 3-OH kinase/AKT and nuclear factor kappa B pathways in human dental pulp stem cells. *Journal of endodontics*, 40(1), 69-75.
- Heidt, T., Courties, G., Dutta, P., Sager, H. B., Sebas, M., Iwamoto, Y., Sun, Y., Da Silva, N., Panizzi, P., & van der Laan, A. M. (2014). Differential Contribution of Monocytes to Heart Macrophages in Steady-State and After Myocardial Infarction Novelty and Significance. *Circulation research*, 115(2), 284-295.
- Heinrichs, D. E., Yethon, J. A., & Whitfield, C. (1998). Molecular basis for structural diversity in the core regions of the lipopolysaccharides of *Escherichia coli* and *Salmonella enterica*. *Molecular microbiology*, 30(2), 221-232.
- Hibbs, J. B. (2002). Infection and nitric oxide. *Journal of Infectious Diseases*, 185(Supplement 1), S9-S17.
- Hollmann, J., & Lindhauer, M. (2005). Pilot-scale isolation of glucuronoarabinoxylans from wheat bran. *Carbohydrate Polymers*, 59(2), 225-230.
- Huang, H., Langenkamp, E., Georganaki, M., Loskog, A., Fuchs, P. F., Dieterich, L. C., Kreuger, J., & Dimberg, A. (2015). VEGF suppresses T-lymphocyte infiltration in the tumor microenvironment through inhibition of NF-κB-induced endothelial activation. *The FASEB Journal*, 29(1), 227-238.
- Hughes, S., Shewry, P., Li, L., Gibson, G., Sanz, M., & Rastall, R. (2007). In vitro fermentation by human fecal microflora of wheat arabinoxylans. *Journal of Agricultural and Food Chemistry*, 55(11), 4589-4595.
- Iqbal, M. S., Akbar, J., Hussain, M. A., Saghir, S., & Sher, M. (2011). Evaluation of hot-water extracted arabinoxylans from ispaghula seeds as drug carriers. *Carbohydrate Polymers*, 83(3), 1218-1225.
- Ise, W., Inoue, T., McLachlan, J. B., Kometani, K., Kubo, M., Okada, T., & Kurosaki, T. (2014). Memory B cells contribute to rapid Bcl6 expression by memory follicular helper T cells. *Proceedings of the National Academy of Sciences*, 111(32), 11792-11797.

- Ito, T., Yamada, A., Batal, I., Yeung, M. Y., McGrath, M. M., Sayegh, M. H., Chandraker, A., & Ueno, T. (2016). the Limits of Linked suppression for regulatory t cells. *Frontiers in immunology*, 7.
- Iwasaki, A., & Pillai, P. S. (2014). Innate immunity to influenza virus infection. *Nature Reviews Immunology*, 14(5), 315-328.
- Iwasaki, A., & Medzhitov, R. (2015). Control of adaptive immunity by the innate immune system. *Nature immunology*, 16(4), 343-353.
- Izydorczyk, M., Biliaderis, C., & Bushuk, W. (1991). Comparison of the structure and composition of water-soluble pentosans from different wheat varieties. *Cereal Chemistry*, 68(2), 139-144.
- Izydorczyk, M., Macri, L., & MacGregor, A. (1998). Structure and physicochemical properties of barley non-starch polysaccharides—I. Water-extractable β -glucans and arabinoxylans. *Carbohydrate Polymers*, 35(3-4), 249-258.
- Izydorczyk, M. S., & Biliaderis, C. G. (1995). Cereal arabinoxylans: advances in structure and physicochemical properties. *Carbohydrate Polymers*, 28(1), 33-48.
- Jakubowska, M. A., Ferdek, P. E., Gerasimenko, O. V., Gerasimenko, J. V., & Petersen, O. H. (2016). Nitric oxide signals are interlinked with calcium signals in normal pancreatic stellate cells upon oxidative stress and inflammation. *Open Biology*, 6(8), 149-160.
- Johnson, G. B., Brunn, G. J., Kodaira, Y., & Platt, J. L. (2002). Receptor-mediated monitoring of tissue well-being via detection of soluble heparan sulfate by Toll-like receptor 4. *The Journal of Immunology*, 168(10), 5233-5239.
- Jones, R. B., Mueller, S., Kumari, S., Vrbanac, V., Genel, S., Tager, A. M., Allen, T. M., Walker, B. D., & Irvine, D. J. (2017). Antigen recognition-triggered drug delivery mediated by nanocapsule-functionalized cytotoxic T-cells. *Biomaterials*, 117, 44-53.
- Kagan, J. C., & Barton, G. M. (2015). Emerging principles governing signal transduction by pattern-recognition receptors. *Cold Spring Harbor perspectives in biology*, 7(3), 153-167.
- Kaminogawa, S., & Nanno, M. (2004). Modulation of immune functions by foods. *Evidence-Based Complementary and Alternative Medicine*, 1(3), 241-250.
- Kanashiro, A., Sônego, F., Ferreira, R. G., Castanheira, F. V., Leite, C. A., Borges, V. F., Nascimento, D. C., Cólón, D. F., Alves-Filho, J. C., & Ulloa, L. (2017). Therapeutic potential and limitations of cholinergic anti-inflammatory pathway in sepsis. *Pharmacological research*, 117, 1-8.
- Kang, H., Lee, M.-G., Lee, J.-K., Choi, Y.-H., & Choi, Y.-S. (2016). Enzymatically-Processed Wheat Bran Enhances Macrophage Activity and Has in Vivo Anti-Inflammatory Effects in Mice. *Nutrients*, 8(4), 188-193.
- Kanjan, P., Sahasrabudhe, N. M., de Haan, B. J., & de Vos, P. (2017). Immune effects of β -glucan are determined by combined effects on Dectin-1, TLR2, 4 and 5. *Journal of Functional Foods*, 37, 433-440.
- Karumuthil-Melethil, S., Gudi, R., Johnson, B. M., Perez, N., & Vasu, C. (2014). Fungal β -glucan, a Dectin-1 ligand, promotes protection from type 1 diabetes by inducing regulatory innate immune response. *The Journal of Immunology*, 193(7), 3308-3321.
- Kawai, T., & Akira, S. (2010). The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nature immunology*, 11(5), 373-384.
- Kim, M. Y., Lee, S. H., Jang, G. Y., Park, H. J., Li, M., Kim, S., Lee, Y. R., Noh, Y. H., Lee, J., & Jeong, H. S. (2015). Effects of high hydrostatic pressure treatment on the enhancement of functional components of germinated rough rice (*Oryza sativa* L.). *Food Chemistry*, 166, 86-92.

- Kim, R., Emi, M., & Tanabe, K. (2006). Functional roles of immature dendritic cells in impaired immunity of solid tumour and their targeted strategies for provoking tumour immunity. *Clinical & Experimental Immunology*, 146(2), 189-196.
- Koegelenberg, D., & Chimphango, A. F. (2017). Effects of wheat-bran arabinoxylan as partial flour replacer on bread properties. *Food Chemistry*, 221, 1606-1613.
- Kozarcanin, H., Lood, C., Munthe-Fog, L., Sandholm, K., Hamad, O. A., Bengtsson, A. A., Skjoedt, M. O., Huber-Lang, M., Garred, P., & Ekdahl, K. N. (2016). The lectin complement pathway serine proteases (MASPs) represent a possible crossroad between the coagulation and complement systems in thromboinflammation. *Journal of Thrombosis and Haemostasis*, 14(3), 531-545.
- Lawrence, T., & Natoli, G. (2011). Transcriptional regulation of macrophage polarization: enabling diversity with identity. *Nature Reviews Immunology*, 11(11), 750-761.
- Le Maitre, C. L. (2014). Role of Cytokines and Proteases in the Pathogenesis of Intervertebral Disc Degeneration. *Global Spine Journal*, 4(S 01), in5. 01.
- Lee, J., Breton, G., Oliveira, T. Y. K., Zhou, Y. J., Aljoufi, A., Pühr, S., Cameron, M. J., Sékaly, R.-P., Nussenzweig, M. C., & Liu, K. (2015). Restricted dendritic cell and monocyte progenitors in human cord blood and bone marrow. *Journal of Experimental Medicine*, 212(3), 385-399.
- Lee, K. Y., Jeong, E. J., Sung, S. H., & Kim, Y. C. (2016). Stemonal alkaloids isolated from *Stemona tuberosa* roots and their inhibitory activity on lipopolysaccharide-induced nitric oxide production. *Records of Natural Products*, 10(1), 109.
- Lerouge, I., & Vanderleyden, J. (2002). O-antigen structural variation: mechanisms and possible roles in animal/plant-microbe interactions. *FEMS microbiology reviews*, 26(1), 17-47.
- Levine, B. (2015). Performance-enhancing drugs: design and production of redirected chimeric antigen receptor (CAR) T cells. *Cancer gene therapy*, 22(2), 79-84.
- Li, W., Zhang, S., & Smith, C. (2015). The molecular structure features-immune stimulatory activity of arabinoxylans derived from the pentosan fraction of wheat flour. *Journal of Cereal Science*, 62, 81-86.
- Li, W., Hu, H., Wang, Q., & Brennan, C. S. (2013). Molecular features of wheat endosperm arabinoxylan inclusion in functional bread. *Foods*, 2(2), 225-237.
- Lim, W. F., Inoue-Yokoo, T., Tan, K. S., Lai, M. I., & Sugiyama, D. (2013). Hematopoietic cell differentiation from embryonic and induced pluripotent stem cells. *Stem cell research & therapy*, 4(3), 71.
- Linehan, E., Dombrowski, Y., Snoddy, R., Fallon, P. G., Kissenpfennig, A., & Fitzgerald, D. C. (2014). Aging impairs peritoneal but not bone marrow-derived macrophage phagocytosis. *Aging cell*, 13(4), 699-708.
- Lovegrove, A., Edwards, C., De Noni, I., Patel, H., El, S., Grassby, T., Zielke, C., Ulmius, M., Nilsson, L., & Butterworth, P. (2017). Role of polysaccharides in food, digestion, and health. *Critical Reviews in Food Science and Nutrition*, 57(2), 237-253.
- Lu, J., Li, Y., Gu, G., & Mao, Z. (2005). Effects of molecular weight and concentration of arabinoxylans on the membrane plugging. *Journal of agricultural and food chemistry*, 53(12), 4996-5002.
- Lu, Q., Yokoyama, C. C., Williams, J. W., Baldrige, M. T., Jin, X., DesRochers, B., Bricker, T., Wilen, C. B., Bagaitkar, J., & Loginicheva, E. (2016). Homeostatic control of innate lung inflammation by Vici syndrome gene *Epg5* and additional autophagy genes promotes influenza pathogenesis. *Cell host & microbe*, 19(1), 102-113.
- Mangoni, M. L., Epand, R. F., Rosenfeld, Y., Peleg, A., Barra, D., Epand, R. M., & Shai, Y. (2008). Lipopolysaccharide, a key molecule involved in the synergism between

- 695 temporins in inhibiting bacterial growth and in endotoxin neutralization. *Journal of*
 696 *Biological Chemistry*, 283(34), 22907-22917.
- 697 Manzo, T., Sturmheit, T., Basso, V., Petrozziello, E., Michelini, R. H., Riba, M., Freschi, M.,
 698 Elia, A. R., Grioni, M., & Curnis, F. (2017). T Cells Redirected to a Minor
 699 Histocompatibility Antigen Instruct Intratumoral TNF α Expression and Empower
 700 Adoptive Cell Therapy for Solid Tumors. *Cancer research*, 77(3), 658-671.
- 701 Marcotuli, I., Hsieh, Y. S.-Y., Lahnstein, J., Yap, K., Burton, R. A., Blanco, A., Fincher, G. B.,
 702 & Gadaleta, A. (2016). Structural Variation and Content of Arabinoxylans in
 703 Endosperm and Bran of Durum Wheat (*Triticum turgidum* L.). *Journal of agricultural*
 704 *and food chemistry*, 64(14), 2883-2892.
- 705 Marletta, M. A. (1994). Nitric oxide synthase: aspects concerning structure and catalysis. *Cell*,
 706 78(6), 927-930.
- 707 Marletta, M. A., Yoon, P. S., Iyengar, R., Leaf, C. D., & Wishnok, J. S. (1988). Macrophage
 708 oxidation of L-arginine to nitrite and nitrate: nitric oxide is an intermediate.
 709 *Biochemistry*, 27(24), 8706-8711.
- 710 Martinez, F. O., & Gordon, S. (2014). The M1 and M2 paradigm of macrophage activation:
 711 time for reassessment. *F1000Prime Rep*, 6(13.10), 12703.
- 712 Martins, V. M., Simões, J., Ferreira, I., Cruz, M. T., Domingues, M. R., & Coimbra, M. A.
 713 (2017). In vitro macrophage nitric oxide production by *Pterospartum tridentatum* (L.)
 714 Willk. inflorescence polysaccharides. *Carbohydrate Polymers*, 157, 176-184.
- 715 Massey, V. L., Poole, L. G., Siow, D. L., Torres, E., Warner, N. L., Schmidt, R. H.,
 716 Ritzenthaler, J. D., Roman, J., & Arteel, G. E. (2015). Chronic Alcohol Exposure
 717 Enhances Lipopolysaccharide-Induced Lung Injury in Mice: Potential Role of Systemic
 718 Tumor Necrosis Factor-Alpha. *Alcoholism: Clinical and Experimental Research*,
 719 39(10), 1978-1988.
- 720 Maude, S. L., Barrett, D., Teachey, D. T., & Grupp, S. A. (2014). Managing cytokine release
 721 syndrome associated with novel T cell-engaging therapies. *Cancer journal (Sudbury,*
 722 *Mass.)*, 20(2), 119.
- 723 McCarthy, C. G., Goulopoulou, S., Wenceslau, C. F., Spitler, K., Matsumoto, T., & Webb, R.
 724 C. (2014). Toll-like receptors and damage-associated molecular patterns: novel links
 725 between inflammation and hypertension. *American Journal of Physiology-Heart and*
 726 *Circulatory Physiology*, 306(2), H184-H196.
- 727 Medzhitov, R. (2001). Toll-like receptors and innate immunity. *Nature Reviews Immunology*,
 728 1(2), 135-145.
- 729 Medzhitov, R., & Janeway Jr, C. (2000). Innate immunity. *New England Journal of Medicine*,
 730 343(5), 338-344.
- 731 Medzhitov, R., Preston-Hurlburt, P., & Janeway, C. A. (1997). A human homologue of the
 732 *Drosophila* Toll protein signals activation of adaptive immunity. *Nature*, 388(6640),
 733 394-397.
- 734 Mendis, M., Leclerc, E., & Simsek, S. (2016). Arabinoxylan hydrolyzates as
 735 immunomodulators in lipopolysaccharide-induced RAW264. 7 macrophages. *Food &*
 736 *Function*, 7(7), 3039-3045.
- 737 Mendis, M., Leclerc, E., & Simsek, S. (2017). Arabinoxylan hydrolyzates as
 738 immunomodulators in Caco-2 and HT-29 colon cancer cell lines. *Food & Function*,
 739 8(1), 220-231.
- 740 Merle, N. S., Church, S. E., Fremeaux-Bacchi, V., & Roumenina, L. T. (2015). Complement
 741 system part I—molecular mechanisms of activation and regulation. *Antimicrobial*
 742 *Peptides and Complement—Maximising the Inflammatory Response*, 17.
- 743 Miller, S. I., Ernst, R. K., & Bader, M. W. (2005). LPS, TLR4 and infectious disease diversity.
 744 *Nature Reviews Microbiology*, 3(1), 36-46.

- Mittal, D., Gubin, M. M., Schreiber, R. D., & Smyth, M. J. (2014). New insights into cancer immunoediting and its three component phases—elimination, equilibrium and escape. *Current opinion in immunology*, 27, 16-25.
- Miwa, T., Sato, S., Gullipalli, D., Nangaku, M., & Song, W.-C. (2013). Blocking properdin, the alternative pathway, and anaphylatoxin receptors ameliorates renal ischemia-reperfusion injury in decay-accelerating factor and CD59 double-knockout mice. *The Journal of Immunology*, 190(7), 3552-3559.
- Moers, K., Celus, I., Brijs, K., Courtin, C. M., & Delcour, J. A. (2005). Endoxylanase substrate selectivity determines degradation of wheat water-extractable and water-unextractable arabinoxylan. *Carbohydrate Research*, 340(7), 1319-1327.
- Mogensen, T. H. (2009). Pathogen recognition and inflammatory signaling in innate immune defenses. *Clinical microbiology reviews*, 22(2), 240-273.
- Mohan, B., Malleshi, N., & Koseki, T. (2010). Physico-chemical characteristics and non-starch polysaccharide contents of Indica and Japonica brown rice and their malts. *LWT-Food Science and Technology*, 43(5), 784-791.
- Mosser, D. M., & Edwards, J. P. (2008). Exploring the full spectrum of macrophage activation. *Nature Reviews Immunology*, 8(12), 958-969.
- Motley, S. T., Morrow, B. J., Liu, X., Dodge, I. L., Vitiello, A., Ward, C. K., & Shaw, K. J. (2004). Simultaneous analysis of host and pathogen interactions during an in vivo infection reveals local induction of host acute phase response proteins, a novel bacterial stress response, and evidence of a host-imposed metal ion limited environment. *Cellular microbiology*, 6(9), 849-865.
- Mucida, D., Husain, M. M., Muroi, S., Van Wijk, F., Shinnakasu, R., Naoe, Y., Reis, B. S., Huang, Y., Lambolez, F., & Docherty, M. (2013). Transcriptional reprogramming of mature CD4⁺ helper T cells generates distinct MHC class II-restricted cytotoxic T lymphocytes. *Nature immunology*, 14(3), 281-289.
- Murofushi, Y., Villena, J., Morie, K., Kanmani, P., Tohno, M., Shimazu, T., Aso, H., Suda, Y., Hashiguchi, K., & Saito, T. (2015). The toll-like receptor family protein RP105/MD1 complex is involved in the immunoregulatory effect of exopolysaccharides from *Lactobacillus plantarum* N14. *Molecular immunology*, 64(1), 63-75.
- Murray, P. J., Allen, J. E., Biswas, S. K., Fisher, E. A., Gilroy, D. W., Goerdts, S., Gordon, S., Hamilton, J. A., Ivashkiv, L. B., & Lawrence, T. (2014). Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity*, 41(1), 14-20.
- Nahrendorf, M., & Swirski, F. K. (2015). Lifestyle effects on hematopoiesis and atherosclerosis. *Circulation research*, 116(5), 884-894.
- Nairz, M., Schleicher, U., Schroll, A., Sonnweber, T., Theurl, I., Ludwiczek, S., Talasz, H., Brandacher, G., Moser, P. L., & Muckenthaler, M. U. (2013). Nitric oxide-mediated regulation of ferroportin-1 controls macrophage iron homeostasis and immune function in *Salmonella* infection. *Journal of Experimental Medicine*, 210(5), 855-873.
- Nicholson, L. B. (2016). The immune system. *Essays in biochemistry*, 60(3), 275-301.
- Nishikawa, H., & Sakaguchi, S. (2014). Regulatory T cells in cancer immunotherapy. *Current opinion in immunology*, 27, 1-7.
- O'Neill, L. A., Golenbock, D., & Bowie, A. G. (2013). The history of Toll-like receptors [mdash] redefining innate immunity. *Nature Reviews Immunology*, 13(6), 453-460.
- O'Neill, L. A., & Pearce, E. J. (2016). Immunometabolism governs dendritic cell and macrophage function. *Journal of Experimental Medicine*, 213(1), 15-23.
- Ogawa, K., Takeuchi, M., & Nakamura, N. (2005). Immunological effects of partially hydrolyzed arabinoxylan from corn husk in mice. *Bioscience, biotechnology, and biochemistry*, 69(1), 19-25.

- 794 Ogawa, M. (1993). Differentiation and proliferation of hematopoietic stem cells. *Blood*, 81(11),
795 2844-2853.
- 796 Ohto, U., Fukase, K., Miyake, K., & Shimizu, T. (2012). Structural basis of species-specific
797 endotoxin sensing by innate immune receptor TLR4/MD-2. *Proceedings of the*
798 *National Academy of Sciences*, 109(19), 7421-7426.
- 799 Okuda, S., Sherman, D. J., Silhavy, T. J., Ruiz, N., & Kahne, D. (2016). Lipopolysaccharide
800 transport and assembly at the outer membrane: the PEZ model. *Nature Reviews*
801 *Microbiology*, 14(12), 337-345.
- 802 Olmos, G., & Lladó, J. (2014). Tumor necrosis factor alpha: a link between neuroinflammation
803 and excitotoxicity. *Mediators of inflammation*, 2014.
- 804 Orlowsky, E. W., & Kraus, V. B. (2015). The role of innate immunity in osteoarthritis: when
805 our first line of defense goes on the offensive. *The Journal of rheumatology*, 42(3), 363-
806 371.
- 807 Orsini, F., De Blasio, D., Zangari, R., Zanier, E. R., & De Simoni, M.-G. (2014). Versatility of
808 the complement system in neuroinflammation, neurodegeneration and brain
809 homeostasis.
- 810 Pacher, P., Beckman, J. S., & Liaudet, L. (2007). Nitric oxide and peroxynitrite in health and
811 disease. *Physiological reviews*, 87(1), 315-424.
- 812 Palygin, O., Endres, B., Miller, B., Sorokin, A., Geurts, A., & Staruschenko, A. (2015). Two-
813 Photon Imaging of Intracellular Ca²⁺ Handling and Nitric Oxide Production in
814 Endothelial and Smooth Muscle Cells of Isolated Rat Vessels. *The FASEB Journal*,
815 29(1), 808-818.
- 816 Park, B. S., Song, D. H., Kim, H. M., Choi, B.-S., Lee, H., & Lee, J.-O. (2009). The structural
817 basis of lipopolysaccharide recognition by the TLR4-MD-2 complex. *Nature*,
818 458(7242), 1191-1195.
- 819 Park, H.-Y., Lee, K.-W., & Choi, H.-D. (2017). Rice bran constituents: immunomodulatory
820 and therapeutic activities. *Food & Function*, 8(3), 935-943.
- 821 Patosuo, T. (2014). Recent research on natural rubber latex (NRL) allergy. *Chem Manuf Appl*
822 *Nat Rubber*, 452-482.
- 823 Pavlovich-Abril, A., Rouzaud-Sáñez, O., Carvajal-Millán, E., Navarro, R. E., Robles-
824 Sánchez, R. M., & Barrón-Hoyos, J. M. (2016). Molecular characterization of water
825 extractable arabinoxylans isolated from wheat fine bran and their effect on dough
826 viscosity. *LWT-Food Science and Technology*, 74, 484-492.
- 827 Peerschke, E. I., Panicker, S., & Bussel, J. (2016). Classical complement pathway activation in
828 immune thrombocytopenia purpura: inhibition by a novel C1s inhibitor. *British journal*
829 *of haematology*, 173(6), 942-945.
- 830 Pietretti, D., Scheer, M., Fink, I. R., Taverne, N., Savelkoul, H. F., Spaink, H. P., Forlenza, M.,
831 & Wiegertjes, G. F. (2014). Identification and functional characterization of
832 nonmammalian Toll-like receptor 20. *Immunogenetics*, 66(2), 123-141.
- 833 Plüddemann, A., Mukhopadhyay, S., & Gordon, S. (2011). Innate immunity to intracellular
834 pathogens: macrophage receptors and responses to microbial entry. *Immunological*
835 *reviews*, 240(1), 11-24.
- 836 Pradere, J.-P., Dapito, D. H., & Schwabe, R. F. (2014). The Yin and Yang of Toll-like receptors
837 in cancer. *Oncogene*, 33(27), 3485.
- 838 Prolo, C., Álvarez, M. N., Ríos, N., Peluffo, G., Radi, R., & Romero, N. (2015). Nitric oxide
839 diffusion to red blood cells limits extracellular, but not intraphagosomal, peroxynitrite
840 formation by macrophages. *Free Radical Biology and Medicine*, 87, 346-355.
- 841 Qiu, S., Yadav, M. P., & Yin, L. (2017). Characterization and functionalities study of
842 hemicellulose and cellulose components isolated from sorghum bran, bagasse and
843 biomass. *Food Chemistry*, 230(3), 225-233.

- Raetz, C. R., Guan, Z., Ingram, B. O., Six, D. A., Song, F., Wang, X., & Zhao, J. (2009). Discovery of new biosynthetic pathways: the lipid A story. *Journal of lipid research*, 50(Supplement), S103-S108.
- Rajilić-Stojanović, M., Jonkers, D. M., Salonen, A., Hanevik, K., Raes, J., Jalanka, J., De Vos, W. M., Manichanh, C., Golic, N., & Enck, P. (2015). Intestinal microbiota and diet in IBS: causes, consequences, or epiphenomena? *The American journal of gastroenterology*, 110(2), 278-287.
- Rath, M., Müller, I., Kropf, P., Closs, E. I., & Munder, M. (2014). Metabolism via arginase or nitric oxide synthase: two competing arginine pathways in macrophages. *Frontiers in immunology*, 5.
- Ren, Q., Lan, J.-F., Zhong, X., Song, X.-J., Ma, F., Hui, K.-M., Wang, W., Yu, X.-Q., & Wang, J.-X. (2014). A novel Toll like receptor with two TIR domains (HcToll-2) is involved in regulation of antimicrobial peptide gene expression of *Hyriopsis cumingii*. *Developmental & Comparative Immunology*, 45(1), 198-208.
- Rietschel, E. T., Kirikae, T., Schade, F. U., Mamat, U., Schmidt, G., Loppnow, H., Ulmer, A. J., Zähringer, U., Seydel, U., & Di Padova, F. (1994). Bacterial endotoxin: molecular relationships of structure to activity and function. *The FASEB Journal*, 8(2), 217-225.
- Rissoan, M.-C., Soumelis, V., Kadowaki, N., Grouard, G., Briere, F., de Waal Malefyt, R., & Liu, Y.-J. (1999). Reciprocal control of T helper cell and dendritic cell differentiation. *Science*, 283(5405), 1183-1186.
- Roubroeks, J., Andersson, R., & Åman, P. (2000). Structural features of (1→3),(1→4)-β-D-glucan and arabinoxylan fractions isolated from rye bran. *Carbohydrate Polymers*, 42(1), 3-11.
- Ryu, J.-K., Kim, S. J., Rah, S.-H., Kang, J. I., Jung, H. E., Lee, D., Lee, H. K., Lee, J.-O., Park, B. S., & Yoon, T.-Y. (2017). Reconstruction of LPS Transfer Cascade Reveals Structural Determinants within LBP, CD14, and TLR4-MD2 for Efficient LPS Recognition and Transfer. *Immunity*, 46(1), 38-50.
- Sadek, M. M., Olia, M. B. A., Nowell, C. J., Barlow, N., Schiesser, C. H., Nicholson, S. E., & Norton, R. S. (2017). Characterisation of a novel coumarin-based fluorescent probe for monitoring nitric oxide production in macrophages. *Bioorganic & Medicinal Chemistry*.
- Sahasrabudhe, N. M., Schols, H. A., Faas, M. M., & Vos, P. (2016). Arabinoxylan activates Dectin-1 and modulates particulate β-glucan-induced Dectin-1 activation. *Molecular nutrition & food research*, 60(2), 458-467.
- Saitoh, S. i., Akashi, S., Yamada, T., Tanimura, N., Kobayashi, M., Konno, K., Matsumoto, F., Fukase, K., Kusumoto, S., & Nagai, Y. (2004). Lipid A antagonist, lipid IVa, is distinct from lipid A in interaction with Toll-like receptor 4 (TLR4)-MD-2 and ligand-induced TLR4 oligomerization. *International immunology*, 16(7), 961-969.
- Sakaguchi, S., Yamaguchi, T., Nomura, T., & Ono, M. (2008). Regulatory T cells and immune tolerance. *Cell*, 133(5), 775-787.
- Sárossy, Z., Tenkanen, M., Pitkänen, L., Bjerre, A.-B., & Plackett, D. (2013). Extraction and chemical characterization of rye arabinoxylan and the effect of β-glucan on the mechanical and barrier properties of cast arabinoxylan films. *Food Hydrocolloids*, 30(1), 206-216.
- Saulnier, L., Sado, P.-E., Branlard, G., Charmet, G., & Guillon, F. (2007). Wheat arabinoxylans: exploiting variation in amount and composition to develop enhanced varieties. *Journal of Cereal Science*, 46(3), 261-281.
- Schneider, C., Nobs, S. P., Kurrer, M., Rehrauer, H., Thiele, C., & Kopf, M. (2014). Induction of the nuclear receptor PPAR-[gamma] by the cytokine GM-CSF is critical for the

- differentiation of fetal monocytes into alveolar macrophages. *Nature immunology*, 15(11), 1026-1037.
- Schnekenburger, M., Dicato, M., & Diederich, M. (2014). Plant-derived epigenetic modulators for cancer treatment and prevention. *Biotechnology Advances*, 32(6), 1123-1132.
- Schwarz, H., Schmittner, M., Duschl, A., & Horejs-Hoeck, J. (2014). Residual endotoxin contaminations in recombinant proteins are sufficient to activate human CD1c+ dendritic cells. *PloS one*, 9(12), e113840.
- Scichilone, N., Battaglia, S., Taormina, S., Modica, V., Pozzecco, E., & Bellia, V. (2013). Alveolar nitric oxide and asthma control in mild untreated asthma. *Journal of Allergy and Clinical Immunology*, 131(6), 1513-1517.
- Silva, M. T. (2010). When two is better than one: macrophages and neutrophils work in concert in innate immunity as complementary and cooperative partners of a myeloid phagocyte system. *Journal of leukocyte biology*, 87(1), 93-106.
- Slocum, C., Coats, S. R., Hua, N., Kramer, C., Papadopoulos, G., Weinberg, E. O., Gudino, C. V., Hamilton, J. A., Darveau, R. P., & Genco, C. A. (2014). Distinct lipid a moieties contribute to pathogen-induced site-specific vascular inflammation. *PLoS Pathog*, 10(7), e1004215.
- Speciale, I., Paciello, I., Fazio, L. L., Sturiale, L., Palmigiano, A., Lanzetta, R., Parrilli, M., Garozzo, D., Lemaitre, B., & Bernardini, M. L. (2015). Determination of the structure of the O-antigen and the lipid A from the entomopathogenic bacterium *Pseudomonas entomophila* lipopolysaccharide along with its immunological properties. *Carbohydrate Research*, 412, 20-27.
- Sperandio, B., Fischer, N., & Sansonetti, P. J. (2015). *Mucosal physical and chemical innate barriers: Lessons from microbial evasion strategies*. Paper presented at the Seminars in immunology.
- Stevenson, M. M., & Riley, E. M. (2004). Innate immunity to malaria. *Nature Reviews Immunology*, 4(3), 169-180.
- Sullivan, C. B., Porter, R. M., Evans, C. H., Ritter, T., Shaw, G., Barry, F., & Murphy, J. M. (2014). TNF α and IL-1 β influence the differentiation and migration of murine MSCs independently of the NF- κ B pathway. *Stem cell research & therapy*, 5(4), 104.
- Takahashi, M., Iwaki, D., Kanno, K., Ishida, Y., Xiong, J., Matsushita, M., Endo, Y., Miura, S., Ishii, N., & Sugamura, K. (2008). Mannose-binding lectin (MBL)-associated serine protease (MASP)-1 contributes to activation of the lectin complement pathway. *The Journal of Immunology*, 180(9), 6132-6138.
- Takeuchi, O., & Akira, S. (2010). Pattern recognition receptors and inflammation. *Cell*, 140(6), 805-820.
- Taylor, P. R., Martinez-Pomares, L., Stacey, M., Lin, H.-H., Brown, G. D., & Gordon, S. (2005). Macrophage receptors and immune recognition. *Annu. Rev. Immunol.*, 23, 901-944.
- Termeer, C., Benedix, F., Sleeman, J., Fieber, C., Voith, U., Ahrens, T., Miyake, K., Freudenberg, M., Galanos, C., & Simon, J. C. (2002). Oligosaccharides of Hyaluronan activate dendritic cells via toll-like receptor 4. *Journal of Experimental Medicine*, 195(1), 99-111.
- Thompson, A. E. (2015). The immune system. *Jama*, 313(16), 1686-1689.
- Trouw, L. A., Pickering, M. C., & Blom, A. M. (2017). The complement system as a potential therapeutic target in rheumatic disease. *Nature reviews. Rheumatology*, 13(9), 538-547.
- Vacchelli, E., Eggermont, A., Sautès-Fridman, C., Galon, J., Zitvogel, L., Kroemer, G., & Galluzzi, L. (2013). Trial Watch: Toll-like receptor agonists for cancer therapy. *Oncoimmunology*, 2(8), e25238.

- van Furth, R., & Cohn, Z. A. (1968). The origin and kinetics of mononuclear phagocytes. *The Journal of experimental medicine*, 128(3), 415.
- Vilček, J., & Feldmann, M. (2004). Historical review: cytokines as therapeutics and targets of therapeutics. *Trends in pharmacological sciences*, 25(4), 201-209.
- Visintin, A., Mazzoni, A., Spitzer, J. H., Wyllie, D. H., Dower, S. K., & Segal, D. M. (2001). Regulation of Toll-like receptors in human monocytes and dendritic cells. *The Journal of Immunology*, 166(1), 249-255.
- Volman, J. J., Ramakers, J. D., & Plat, J. (2008). Dietary modulation of immune function by β -glucans. *Physiology & behavior*, 94(2), 276-284.
- Wang, J., Sun, B., Liu, Y., & Zhang, H. (2014). Optimisation of ultrasound-assisted enzymatic extraction of arabinoxylan from wheat bran. *Food Chemistry*, 150, 482-488.
- Wang, J., Smits, E., Boom, R. M., & Schutyser, M. A. (2015). Arabinoxylans concentrates from wheat bran by electrostatic separation. *Journal of Food Engineering*, 155, 29-36.
- Wiersinga, W. J., Leopold, S. J., Cranendonk, D. R., & van der Poll, T. (2014). Host innate immune responses to sepsis. *Virulence*, 5(1), 36-44.
- Witztum, J. L., & Lichtman, A. H. (2014). The influence of innate and adaptive immune responses on atherosclerosis. *Annual Review of Pathology: Mechanisms of Disease*, 9, 73-102.
- Wong, G., Kobinger, G. P., & Qiu, X. (2014). Characterization of host immune responses in Ebola virus infections. *Expert review of clinical immunology*, 10(6), 781-790.
- Wu, E. L., Engström, O., Jo, S., Stuhlsatz, D., Yeom, M. S., Klauda, J. B., Widmalm, G., & Im, W. (2013). Molecular dynamics and NMR spectroscopy studies of E. coli lipopolysaccharide structure and dynamics. *Biophysical journal*, 105(6), 1444-1455.
- Yu, L., Yakubov, G. E., Zeng, W., Xing, X., Stenson, J., Bulone, V., & Stokes, J. R. (2017). Multi-layer mucilage of *Plantago ovata* seeds: Rheological differences arise from variations in arabinoxylan side chains. *Carbohydrate Polymers*, 165, 132-141.
- Yu, M., Shi, W., Zhang, J., Niu, L., Chen, Q., Yan, D., Liu, T., Jing, W., Jiang, X., & Wei, F. (2009). Influence of reverse signaling via membrane TNF- α on cytotoxicity of NK92 cells. *European journal of cell biology*, 88(3), 181-191.
- Zeng, Y.-Y., Jiang, W.-D., Liu, Y., Wu, P., Zhao, J., Jiang, J., Kuang, S.-Y., Tang, L., Tang, W.-N., & Zhang, Y.-A. (2016). Dietary alpha-linolenic acid/linoleic acid ratios modulate intestinal immunity, tight junctions, anti-oxidant status and mRNA levels of NF- κ B p65, MLCK and Nrf2 in juvenile grass carp (*Ctenopharyngodon idella*). *Fish & shellfish immunology*, 51, 351-364.
- Zhang, S., Li, W., Smith, C., & Musa, H. (2015). Cereal-derived arabinoxylans as biological response modifiers: extraction, molecular features, and immune-stimulating properties. *Critical Reviews in Food Science and Nutrition*, 55(8), 1035-1052.
- Zhang, Z., Smith, C., Li, W., & Ashworth, J. (2016). Characterization of nitric oxide modulatory activities of alkaline-extracted and enzymatic-modified arabinoxylans from corn bran in cultured human monocytes. *Journal of Agricultural and Food Chemistry*, 64(43), 8128-8137.
- Zhou, S., Liu, X., Guo, Y., Wang, Q., Peng, D., & Cao, L. (2010). Comparison of the immunological activities of arabinoxylans from wheat bran with alkali and xylanase-aided extraction. *Carbohydrate Polymers*, 81(4), 784-789.
- Zhu, E. F., Gai, S. A., Opel, C. F., Kwan, B. H., Surana, R., Mihm, M. C., Kauke, M. J., Moynihan, K. D., Angelini, A., & Williams, R. T. (2015). Synergistic innate and adaptive immune response to combination immunotherapy with anti-tumor antigen antibodies and extended serum half-life IL-2. *Cancer cell*, 27(4), 489-501.

991 Table 1. Water extractable and water-unextractable AXs in some cereals (dry weight basis)
 992 weight basis

WEAX and WUAX in some cereal grains and cereal by-products (dry weight basis)					
Cereal	Tissues	Total AXs %	WEAX %	WUAX %	References
Rice	Bran	4.84-5.11	0.35-0.77	4.34-4.49	(Hashimoto, Shogren, Bolte, et al., 1987)
	Bran	8.5	0.2	8.3	(Choct, 1997)
	Hulls	8.36-9.24	0.11	8.25-9.13	(Hashimoto, Shogren, Bolte, et al., 1987)
	Cooked	0.5	NA	NA	(Dodevska et al., 2013)
	Germinated whole grain	2.97-6.84	NA	NA	(Kim et al., 2015)
	Whole grain	2.64	0.06	2.58	(Hashimoto, Shogren, Bolte, et al., 1987)
Wheat	Bran	25	1	24	(Hollmann & Lindhauer, 2005)
	De-starched bran	29.1	NA	NA	(Koegelenberg & Chiphango, 2017)
	Bran	26.2	NA	NA	(Koegelenberg & Chiphango, 2017)
	Bran	23	NA	NA	(Wang et al., 2015)
	Bran	19.38	0.88	18.5	(Hashimoto, Shogren, & Pomeranz, 1987)
	Endosperm	NA	8.23	NA	(Li et al., 2015)
	Endosperm	1.5-2.5	0.3-0.75	1.2-1.7	(Li et al., 2013)
	Endosperm	1.52-1.75	0.42-0.68	1.07-1.1	(Marcotuli et al., 2016)
	Flour	1.37-2.06	0.54-0.68	0.83-1.38	(Izydorczyk et al., 1991)
	White flour	5.1	2.1	2.96	(Pavlovich-Abril et al., 2016)
	Whole grain	5.77	0.59	5.18	(Hashimoto, Shogren, & Pomeranz, 1987)
	Whole grain	8.1	1.8	6.3	(Choct, 1997)

Rye	Whole grain	8-12.1	2.6-4.1	5.4-8	(Hansen et al., 2003)
	Bran	13	2.86-4.29	8.71-10.14	(Sárossy et al., 2013)
	Flour	3.2-3.64	2.2-2.65	0.99-1	(Cyran et al., 2003)
	Whole grain	8.9	3.4	5.5	(Choct, 1997)
Corn	Bran	29.86	0.28	29.58	(Hashimoto, Shogren, Bolte, et al., 1987)
	Bran	26.0	0.71	25.29	(Zhang et al., 2016)

993

994

Table2. Structural properties of AXs

Origin	Extraction method	Immunomodulatory activity	Mw (kDa)	Glu %	Gal %	Xyl %	Ara %	Ar/Xy	References
Wheat bran	alkaline	Tumour inhibition, Mφ activation	352	7.7	na	50.2	41.8	0.83	(Zhou et al., 2010)
Wheat bran	enzyme	Mφ activation	32.5	2.8	na	62.4	34.8	0.55	(Zhou et al., 2010)
Rice bran	enzyme	Mφ, DCs and NK activation	30-50	6	5-7	48-54	22-26	0.5	(Zhang et al., 2015)

998 **Figure Legends**

999 Figure 1. Simplified overview of the immune system; IL- interleukin, TNF α - tumour necrosis
1000 factor α , IF γ – interferon γ . Adapted from (Dranoff, 2004; Stevenson & Riley, 2004).

1001 Figure 2. Lipopolysaccharide (LPS) structure

1002 a. LPS lipid A, O-antigen and core oligosaccharide. b. TLR4-MD2-CD14 receptor
1003 complex (Miller et al., 2005).


1004 Figure 3. Arabinoxylan structure (Garófalo et al., 2011).

1005 Figure 4. Simplified schematic representation of arabinoxylans AXs

1006 (a) Wheat flour and (b) Wheat bran. Substituents above and below the backbone represent
1007 C (O)-2 and C (O)-3 positions, respectively (Edwards et al., 2003; Zhou et al., 2010;
1008 Qiu et al., 2017).


1009 Figure 5. Simplified representation of the macrophage TLR 4 receptor with AXs and LPS

1010



1011

1012 FIGURE 1. Simplified overview of the immune system; IL- interleukin, TNF α - tumour
 1013 necrosis factor α , IF γ – interferon γ . Adapted from (Dranoff, 2004; Stevenson & Riley, 2004).



1014

1015 **FIGURE 2.** Lipopolysaccharide (LPS) structure

1016 a. LPS lipid A, O-antigen and core oligosaccharide. b. TLR4-MD2-CD14 receptor complex. Source. Miller et al.
 1017 (2005).

1018




FIGURE 3 Arabinoxylan structure. Source. Garófalo, Vazquez , Ferreira and Soule (2011).




a**b**

FIGURE 4 Simplified schematic representation of arabinoxylans AXs. (a) Wheat flour and (b) Wheat bran. Substituents above and below the backbone represent C (O)-2 and C (O)-3 positions, respectively. Source. Edwards, Chaplin, Blackwood, and Dettmar (2003), Qiu et al. (2017), and Zhou et al. (2010)




FIGURE 5. Simplified representation of the monocyte TLR 4 receptor with AXs and LPS